


IMAGES IN CARDIOLOGY

Cardiac Allograft Vasculopathy Progression Associated With Intraplaque Neovascularization

Yasuhiro Ichibori, MD,* Daisaku Nakatani, MD, PhD,* Yasushi Sakata, MD, PhD,* Kouichi Tachibana, MD, PhD,* Takashi Akasaka, MD, PhD,† Shunsuke Saito, MD, PhD,‡ Norihide Fukushima, MD, PhD,‡ Yoshiki Sawa, MD, PhD,‡ Shinsuke Nanto, MD, PhD,* Issei Komuro, MD, PhD*

Osaka and Wakayama, Japan



From the *Department of Cardiovascular Medicine, Osaka University Graduate School of Medicine, Osaka, Japan; †Department of Cardiovascular Medicine, Wakayama Medical University, Wakayama, Japan; and the ‡Department of Cardiovascular Surgery, Osaka University Graduate School of Medicine, Osaka, Japan. Dr. Akasaka has received research support from Abbott Vascular Japan, Boston Scientific Japan, Goodman Inc., and St. Jude Medical Japan, and Terumo Inc. and consulting fees from Goodman Inc., St. Jude Medical Japan, and Terumo Inc. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Manuscript received August 17, 2012; accepted August 26, 2012.

A 52-year-old man underwent scheduled cardiac catheterization for assessment of cardiac allograft vasculopathy (CAV) at 3 and 12 months after heart transplantation (HTx). Intravascular ultrasound showed only mild CAV in the left anterior descending coronary artery (LAD) at 3 months (**B**, [Online Video 1](#)). However, at 12 months, CAV progression was detected, predominantly in the proximal LAD (**A and C3**, [Online Video 2](#)). Corresponding optical coherence tomography images revealed the presence of many no-signal tubuloluminal structures within the fibrous plaque in serial frames. Parts of those structures were found to be connected to the vessel lumen, indicating neovascularization (**D2, D3, D4, and E [arrows]**, [Online Video 3](#)). No neovascularization was observed in lesions without CAV progression (**C1 and D1**). To our best knowledge, this is the first reported case demonstrating CAV progression associated with intraplaque neovascularization in vivo. Intraplaque neovascularization could play an important role in CAV progression. DB = diagonal branch; LCX = left circumflex coronary artery.